

Aripiprazole

Quick Reference Guide for Teachers

Aripiprazole is used to treat mood disorders such as bipolar disorder, schizophrenia, and irritability associated with autistic disorder. It is used in combination with other medication to treat depression.

Possible Side Effects	<ul style="list-style-type: none"> • Dizziness • Drowsiness • Vomiting • Drooling • Blurred vision • Weight gain • Constipation 	<ul style="list-style-type: none"> • Fast/pounding heartbeat • Fainting • Mood changes (e.g. anxiety, depression) • Trouble swallowing 	<ul style="list-style-type: none"> • Restlessness • Shaking (tremor) • Muscle spasm/stiffness • Mask-like facial expressions • Seizures
	Signs of infection: Fever, Persistent sore throat		

When To Get Help	<p>Seek immediate medical help if you see: Symptoms of a serious allergic reaction:</p> <ul style="list-style-type: none"> • Rash • Itching/swelling (of face/tongue/throat) • Severe dizziness • Trouble breathing
	<p>In case of overdose, contact the local poison centre or emergency room immediately. Symptoms of overdose include:</p> <ul style="list-style-type: none"> • Very fast heartbeat • Loss of consciousness <p>In rare cases, Aripiprazole can cause a condition called Tardive Dyskinesia. Symptoms include:</p> <ul style="list-style-type: none"> • Unusual uncontrolled movements of the face, mouth, tongue, arms, or legs <p>Aripiprazole can also cause a condition called Neuroleptic Malignant Syndrome. Symptoms include:</p> <ul style="list-style-type: none"> • Fever • Muscle stiffness • Severe confusion • Sweating • Fast/irregular heartbeat

Aripiprazole

Quick Reference Guide for Teachers

Aripiprazole is used to treat mood disorders such as bipolar disorder, schizophrenia, and irritability associated with autistic disorder. It is used in combination with other medication to treat depression.

Possible Side Effects	<ul style="list-style-type: none"> • Dizziness • Drowsiness • Vomiting • Drooling • Blurred vision • Weight gain • Constipation 	<ul style="list-style-type: none"> • Fast/pounding heartbeat • Fainting • Mood changes (e.g. anxiety, depression) • Trouble swallowing 	<ul style="list-style-type: none"> • Restlessness • Shaking (tremor) • Muscle spasm/stiffness • Mask-like facial expressions • Seizures
	Signs of infection: Fever, Persistent sore throat		

When To Get Help	<p>Seek immediate medical help if you see: Symptoms of a serious allergic reaction:</p> <ul style="list-style-type: none"> • Rash • Itching/swelling (of face/tongue/throat) • Severe dizziness • Trouble breathing
	<p>In case of overdose, contact the local poison centre or emergency room immediately. Symptoms of overdose include:</p> <ul style="list-style-type: none"> • Very fast heartbeat • Loss of consciousness <p>In rare cases, Aripiprazole can cause a condition called Tardive Dyskinesia. Symptoms include:</p> <ul style="list-style-type: none"> • Unusual uncontrolled movements of the face, mouth, tongue, arms, or legs <p>Aripiprazole can also cause a condition called Neuroleptic Malignant Syndrome. Symptoms include:</p> <ul style="list-style-type: none"> • Fever • Muscle stiffness • Severe confusion • Sweating • Fast/irregular heartbeat