

# Risperidone

## Quick Reference Guide for Teachers

Risperidone, also known as Risperidal, is used to treat psychosis in individuals who have a psychotic illness.

- |                              |   |   |
|------------------------------|---|---|
| <b>Possible Side Effects</b> | <ul style="list-style-type: none"> <li>• Sleepiness, fatigue</li> <li>• Stomach discomfort</li> <li>• Nausea</li> <li>• Urine incontinence</li> <li>• Increased saliva</li> <li>• Dizziness</li> <li>• Spontaneous flow of breast milk</li> </ul> | <ul style="list-style-type: none"> <li>• Weight gain</li> <li>• Slow, absent movements</li> <li>• Muscle stiffness or rigidity</li> <li>• Slow shuffling walk</li> <li>• Drooling</li> <li>• Restlessness</li> <li>• Fidgeting</li> </ul> |
|------------------------------|---|---|

<b>When To Get Help</b>	<p><b>Seek immediate medical help if you see:</b></p> <p>Symptoms of Tardive dyskinesia which involve abnormal involuntary movements:</p> <ul style="list-style-type: none"> <li>• Lip-smacking</li> <li>• Tongue-thrusting</li> <li>• Slow writhing</li> </ul> <p>Symptoms of Acute Dystonia, a rapid, painful muscular spasm which can take the form of:</p> <ul style="list-style-type: none"> <li>• Neck muscle contraction</li> <li>• Extreme rolling back of the eyes with rapid blinking</li> <li>• Arching of the neck and back.</li> </ul> <p>Symptoms of Neuroleptic Malignant Syndrome:</p> <ul style="list-style-type: none"> <li>• Fever</li> <li>• Incontinence</li> <li>• Muscle rigidity</li> <li>• An altered level of consciousness (i.e. disorientation, reduced alertness)</li> </ul>
-------------------------	---

# Risperidone

## Quick Reference Guide for Teachers

Risperidone, also known as Risperidal, is used to treat psychosis in individuals who have a psychotic illness.

- |                              |   |   |
|------------------------------|---|---|
| <b>Possible Side Effects</b> | <ul style="list-style-type: none"> <li>• Sleepiness, fatigue</li> <li>• Stomach discomfort</li> <li>• Nausea</li> <li>• Urine incontinence</li> <li>• Increased saliva</li> <li>• Dizziness</li> <li>• Spontaneous flow of breast milk</li> </ul> | <ul style="list-style-type: none"> <li>• Weight gain</li> <li>• Slow, absent movements</li> <li>• Muscle stiffness or rigidity</li> <li>• Slow shuffling walk</li> <li>• Drooling</li> <li>• Restlessness</li> <li>• Fidgeting</li> </ul> |
|------------------------------|---|---|

<b>When To Get Help</b>	<p><b>Seek immediate medical help if you see:</b></p> <p>Symptoms of Tardive dyskinesia which involve abnormal involuntary movements:</p> <ul style="list-style-type: none"> <li>• Lip-smacking</li> <li>• Tongue-thrusting</li> <li>• Slow writhing</li> </ul> <p>Symptoms of Acute Dystonia, a rapid, painful muscular spasm which can take the form of:</p> <ul style="list-style-type: none"> <li>• Neck muscle contraction</li> <li>• Extreme rolling back of the eyes with rapid blinking</li> <li>• Arching of the neck and back.</li> </ul> <p>Symptoms of Neuroleptic Malignant Syndrome:</p> <ul style="list-style-type: none"> <li>• Fever</li> <li>• Incontinence</li> <li>• Muscle rigidity</li> <li>• An altered level of consciousness (i.e. disorientation, reduced alertness)</li> </ul>
-------------------------	---

This reference guide was supported from the Canadian Institutes of Health Research (FRN: KAL-104248). Visit <http://katydid.info> for more information.



This reference guide was supported from the Canadian Institutes of Health Research (FRN: KAL-104248). Visit <http://katydid.info> for more information.

